

Hemolymph Acid–Base Balance of Akoya Pearl Oyster *Pinctada fucata martensii* with Cannulated Adductor Muscle in Normoxic Conditions

HANDA Takeshi^{1 †}, ARAKI Akira¹ and YAMAMOTO Ken-ichi¹

Abstract : We collected the hemolymph anaerobically through a cannula inserted into the adductor muscle of Akoya pearl oyster *Pinctada fucata martensii* submerged in experimental seawater, and examined hemolymph pH, total CO₂ concentration (Tco₂), CO₂ partial pressure (Pco₂), and bicarbonate concentration ([HCO₃⁻]) in order to evaluate the ability of the acid–base balance of *P. fucata martensii* in normoxic conditions. The mean values of hemolymph pH and Tco₂ were 7.544 and 2.21 mM/L, respectively. The apparent dissociation constant of carbonic acid (pKapp) was estimated using the following equation: $pKapp = 225.712 - 97.096 \cdot pH + 14.313 \cdot pH^2 - 0.7035 \cdot pH^3$. Using mean values of pKapp (5.99585) and CO₂ solubility coefficient (α_{CO_2} , 37.02 μ M/L/torr) determined in this study, the hemolymph Pco₂ and [HCO₃⁻] were calculated as 1.69 torr and 2.14 mM/L, respectively. The non-bicarbonate buffer value (β_{NB}) was 0.967 slykes.

Key words : *Pinctada fucata martensii*, Akoya pearl oyster, hemolymph, acid–base balance, adductor muscle, buffer value

Introduction

Akoya pearl oyster *Pinctada fucata martensii* is a filibranchial bivalve classified in the Pteriidae and is endemic to Japan.¹⁾ *P. fucata martensii* is distributed from the Boso Peninsula on the Pacific coast and the Oga Peninsula on the Japan Sea coast south to Okinawa.¹⁾ *P. fucata martensii* has nacreous aragonite in the inner layer of its shell valves, and it is used for the production of Akoya pearls. The process of pearl production is similar to the growth of the shell valves and is directly related to metabolism. The metabolism of *P. fucata martensii* has been studied in terms of regulation of gill ventilation volume and oxygen uptake in normoxic, hypoxic, anathermal, and feeding conditions.²⁻⁶⁾ The anatomical structures of the digestive diverticula, ctenidium, and circulatory system were clarified recently.^{7,8)} Handa and Yamamoto (2011, 2012) developed surgical procedures, cannulation of the anterior aorta in *P. fucata martensii*, and examined the hemolymph oxygen and acid–base status in the anterior aorta.^{9,10)} In this

study, we developed cannulation of the adductor muscle in Akoya pearl oyster and examined its hemolymph properties, including acid–base balance. Research into the acid–base balance may contribute to efficient CO₂ utilization, which is related to respiration and calcification for the formation of the shell valves. The CO₂ dynamic phase of *P. fucata martensii* is useful for evaluation of the cultivation environments, and of the effects of ocean acidification and increases in CO₂ level. In some marine bivalves, the CO₂ partial pressure (Pco₂) of the hemolymph is 0.57–2.3 torr in normoxic and normocapnic conditions.¹¹⁻¹⁷⁾ The estimation CO₂ partial pressure by application of the Henderson–Hasselbalch equation is practiced in studies of acid–base balance owing to the relative ease and accuracy of such estimates.¹⁸⁾ In the equation, the characteristic values of the CO₂ solubility coefficient (α_{CO_2}) and apparent dissociation constant of carbonic acid (pKapp) in the hemolymph are required. Therefore, we determined hemolymph α_{CO_2} and pKapp of *P. fucata martensii*, and evaluated the hemolymph acid–base balance in comparison with the other bivalves.

Materials and Methods

Experimental animals and conditions

The experiments used 16 Akoya pearl oysters *P. fucata martensii* (total wet weight: 42.6 ± 2.6 g (mean \pm SE)). The animals were obtained from a marine farm in Tsushima, Nagasaki Prefecture, Japan. After cleaning the shell valves, they were reared for 1 month at 26°C in aerated seawater with added cultivated phytoplankton.^{3,19,20} Twenty-four hours before collecting hemolymph, *P. fucata martensii* was transferred to particle-free (>0.45 μ m) seawater. All experiments were conducted in seawater with a salinity of 33 psu, water temperature 26°C, O₂ saturation 100%, pH 8.19, and total CO₂ concentration 1.7 mM/L.

Surgical procedures and hemolymph collection

Hemolymph was collected from the adductor muscle using a cannula (PE-50; Clay-Adams polyethylene tubing, 0.96 mm outer diameter, 0.58 mm inner diameter, Becton Dickinson Co., US). A small incision was made in the shell valve from the center of the posterior margin. A cannula with a stylet was inserted through the incision into the adductor muscle and advanced toward the center of the adductor muscle. The stylet was removed, and the outside of the cannula was closed. The cannula was gently fixed to the left shell valve using denture adhesive (Kobayashi Pharmaceutical Co., JP) in order to prevent effects from movement of the shell valves. The cannulated oyster was transferred to the acrylic respiratory chamber and was allowed to recover for 2–3 h at 26.0 ± 0.2 °C in normoxic conditions. A hemolymph sample was then drawn through the cannula using a gas-tight micro syringe (1750LTN; Hamilton Co., US). The volume of hemolymph collected was 0.3–0.4 mL.

Hemolymph analysis

Hemolymph pH and total CO₂ concentration (Tco₂) were measured immediately after each collection. The pH was measured using a blood gas meter (BGM200; Cameron Instruments Co., US) with glass and reference electrodes (E301, E351; Cameron Instruments Co., US) at

26°C. Tco₂ was measured using a total CO₂ analyzer (Capnicon 5; Cameron Instruments Co., US). Hemolymph total protein concentration (Tpro, g/L) was determined with a test kit (Micro TP-test; Wako Pure Chemical Co., JP) and a spectrophotometer (Spectronic 20A; Shimadzu Co., JP).

The hemolymph Pco₂ and bicarbonate concentration (HCO₃⁻) were calculated by rearranging the Henderson–Hasselbalch equation.^{21,22} In the equation, α co₂ and pKapp of *P. fucata martensii* were required. The determinations of α co₂ and pKapp were performed using *in vitro* experiments. The α co₂ was determined using hemolymph, which was adjusted to pH 2.5 by the addition of lactic acid (Wako Pure Chemical Co., JP). The acidified sample was transferred to a tonometer flask and equilibrated with humidified standard CO₂ gas (CO₂, 10.0%; O₂, 20.9%; N₂ balance) using an equilibrator (DEQ-1; Cameron Instruments Co., US) at 26°C, and subsequently the Tco₂ of each equilibrated sample was measured using a total CO₂ analyzer. The Pco₂ of the equilibrated sample was calculated from known CO₂ concentration standard gas (10.0%), prevailing barometric pressure, and water vapor pressure at 26°C. For the determination of pKapp, the hemolymph sample was transferred to a tonometer flask and equilibrated with humidified standard CO₂ gases (CO₂, 0.1%, 0.2%, 0.5%, 1.0%, and 2.0%; O₂, 20.9%; N₂ balance) using an equilibrator at 26°C. After equilibration, the pH and Tco₂ of the sample were measured using a blood gas meter and total CO₂ analyzer. Using the sample pH, Tco₂, and α co₂, pKapp was determined by rearrangement of Henderson–Hasselbalch equation.^{18,21}

Calculation

α co₂ of the experimental animals was calculated using the equation:

$$\alpha\text{co}_2 = \text{Tco}_2 \cdot \text{Pco}_2^{-1}$$

where the units of the parameters are mM/L/torr for α co₂, mM/L for Tco₂ and torr for Pco₂.

Using the sample pH, Tco₂, and α co₂ calculated using the above equation, pKapp was determined by

rearrangement of Henderson–Hasselbalch equation^{18,21)} as follows:

$$\text{pKapp} = \text{pH} - \log [(T\text{co}_2 - \alpha\text{co}_2 \cdot \text{Pco}_2) \cdot (\alpha\text{co}_2 \cdot \text{Pco}_2)^{-1}]$$

where Pco_2 is calculated from known CO_2 concentration standard gases.

The hemolymph Pco_2 and $[\text{HCO}_3^-]$ were calculated using the equations:

$$\begin{aligned} \text{Pco}_2 &= T\text{co}_2 \cdot [\alpha\text{co}_2 \cdot (1 + 10^{(\text{pH} - \text{pKapp})})]^{-1} \\ [\text{HCO}_3^-] &= T\text{co}_2 - \alpha\text{co}_2 \cdot \text{Pco}_2 \end{aligned}$$

where $T\text{co}_2$ and pH were measured values, and αco_2 and pKapp were obtained in *in vitro* experiments.

The non-bicarbonate buffer value (β_{NB} , slykes) was obtained as the regression coefficient relating pH and $[\text{HCO}_3^-]$ from the *in vitro* experiment.

Statistical analysis

All data are expressed as means \pm standard error. Kruskal–Wallis test was performed for changes in pH and pKapp using the CO_2 standard gases. The comparison of two parameters used Mann–Whitney *U* test. Statistically significant differences were set at $P < 0.05$ (KyPlot 5.0, KyensLab Inc., JP).

Results

Hemolymph samples were collected anaerobically through a cannula from the adductor muscle of *P. fucata martensii*. The mean values of hemolymph pH and $T\text{co}_2$ in normoxic conditions were 7.544, and 2.21 mM/L, respectively (Table 1). The hemolymph αco_2 was 37.02 $\mu\text{M/L/torr}$. The hemolymph pKapp at known CO_2 partial pressures (standard gases) and the corresponding measured pH and $T\text{co}_2$ values are shown in Table 2. The calculated pKapp from all hemolymph samples was 5.995853 ± 0.051311 . Hemolymph Pco_2 and $[\text{HCO}_3^-]$ were

Table 1. Hemolymph pH , total CO_2 concentration ($T\text{co}_2$), CO_2 partial pressure (Pco_2) and bicarbonate concentration ($[\text{HCO}_3^-]$) of the Akoya pearl oyster *Pinctada fucata martensii* at 26°C under normoxic condition

		Mean	SE	N
pH		7.544	0.039	8
$T\text{co}_2$	mM/L	2.21	0.066	8
Pco_2	torr	1.69	0.164	8
$[\text{HCO}_3^-]$	mM/L	2.14	0.062	8

αco_2 : 37.02 $\mu\text{M/L/torr}$; pKapp : 5.995853 (Mean values)

Table 2. Mean values of measured pH , total CO_2 concentration ($T\text{co}_2$), and calculated apparent dissociation constant of carbonic acid (pKapp) of the hemolymph in adductor muscle of the Akoya pearl oyster *Pinctada fucata martensii* with known Pco_2 standard gases

Standard gas		Hemolymph			
CO_2 (%)	Pco_2 (torr)	pH	$T\text{co}_2$ (mM/L)	pKapp	N
0.102	0.75	7.521	1.478	5.80686	8
0.203	1.49	7.389	1.654	5.91215	8
0.515	3.78	7.185	1.977	6.06691	8
1.01	7.41	6.964	2.356	6.08460	8
2.00	14.7	6.714	2.750	6.10875	8

Water temperature 26.0°C

calculated by substitution of the mean value of a_{CO_2} and pK_{app} in the rearranged Henderson-Hasselbalch equation as follows:

$$P_{CO_2} = T_{CO_2} \cdot [0.03702 \cdot (1 + 10^{(pH - 5.995853)})]^{-1}$$

$$[HCO_3^-] = T_{CO_2} - 0.03702 \cdot P_{CO_2}$$




Fig. 1. Relationship between pH and apparent dissociation constant of carbonic acid (pK_{app}) of hemolymph collected from the adductor muscle of Akoya pearl oyster *Pinctada fucata martensii* at 26°C. Values are means \pm standard error. Solid curve fitted to the data and the equation: $pK_{app} = 225.712 - 97.096 \cdot pH + 14.313 \cdot pH^2 - 0.7035 \cdot pH^3$ ($R^2 = 0.9871$)

where the units of the parameters in the equations are torr for P_{CO_2} and mM/L for T_{CO_2} and $[HCO_3^-]$.

Hemolymph P_{CO_2} and $[HCO_3^-]$ at 26°C in normoxic conditions were 1.69 torr and 2.14 mM/L, respectively (Table 1). In *in vitro* experiments (Table 2), the changes in pH and pK_{app} were statistically significant with the increase in P_{CO_2} ($P < 0.05$). At the same time, the interaction between pK_{app} and pH was analyzed (Fig. 1), and the correction equation for pK_{app} was obtained as follows:

$$pK_{app} = 225.712 - 97.096 \cdot pH + 14.313 \cdot pH^2 - 0.7035 \cdot pH^3$$

For comparison, P_{CO_2} and $[HCO_3^-]$ were estimated using the mean value of pK_{app} and the correction equation (Table 3). The hemolymph P_{CO_2} estimated by the correction equation was lower than that from the mean value of pK_{app} ($P < 0.05$). The non-bicarbonate buffer value (β_{NB}), which was obtained as the regression coefficient relating pH and $[HCO_3^-]$ in the *in vitro* experiment, was 0.967 slykes (Table 4). The hemolymph T_{pro} was 1.6 ± 0.04 g/L.

Table 3. The comparison of the values calculated by the correction equation and by the mean pK_{app} in hemolymph P_{CO_2} and $[HCO_3^-]$

	P_{CO_2} (torr)	$[HCO_3^-]$ (mM/L)	N
the mean value of pK_{app}	1.69	2.14	8
pK_{app} calculated using the correction equation	1.06 *	2.17	8

* : statistically significant difference (Mann-Whitney U test, $P < 0.05$)

Table 4. Mean values of measured pH and calculated bicarbonate concentration ($[HCO_3^-]$) of the hemolymph of Akoya pearl oyster *Pinctada fucata martensii* with known P_{CO_2} standard gases

Standard gas		Hemolymph		
CO_2 (%)	P_{CO_2} (torr)	pH	$[HCO_3^-]$ (mM/L)	N
0.102	0.750	7.521	1.45	8
0.203	1.49	7.389	1.60	8
0.515	3.78	7.185	1.84	8
1.01	7.41	6.964	2.08	8
2.00	14.7	6.714	2.21	8

Water temperature: 26.0°C; Non-bicarbonate buffer value (β_{NB}): 0.967

Discussion

We collected the hemolymph of *P. fucata martensii* anaerobically through a cannula from submerged experimental animals after pretreatment by adductor muscle catheterization, and examined hemolymph properties in order to evaluate the acid–base balance of *P. fucata martensii* in normoxic conditions.

P. fucata martensii hemolymph in the adductor muscle exhibited pH of 7.544 and Tco₂ of 2.21 mM/L at 26.0°C. Previously reported hemolymph pH values include pH 7.65 in *Mytilus edulis* at 12°C,¹¹ pH 7.55 in *M. galloprovincialis* at 18°C,¹² pH 7.414 in *Crassostrea gigas* at 23°C,¹⁶ pH 7.617 in *M. coruscus* at 24°C,¹⁵ pH 7.442 in noble scallop *Mimachlamys nobilis* at 24°C,¹⁴ pH 7.576 in *Ostrea denselamellosa* at 24°C,¹⁷ and pH 7.563 in *P. margaritifera* at 26°C.¹³ The hemolymph pH in *P. fucata martensii* was almost the same as that in *M. galloprovincialis*, *O. denselamellosa*, and *P. margaritifera*. Handa and Yamamoto (2015, 2016) and Handa et al. (2017, 2018) reported hemolymph Tco₂ in *P. margaritifera*, *C. gigas*, *M. nobilis*, *M. coruscus*, and *O. denselamellosa* as 2.04 mM/L, 1.87 mM/L, 1.50 mM/L, 1.44 mM/L, and 1.22 mM/L, respectively.^{13–17} The content of carbonic acid and CO₂ was approximately the same as *P. margaritifera*, and higher than that in *C. gigas*, *M. nobilis*, *M. coruscus*, and *O. denselamellosa*. *P. fucata martensii* had an acid–base status that was similar to *P. margaritifera* (Pteriidae bivalve), but seemed to be different from Mytilidae, Pectinidae, and Ostreidae bivalves.

Cameron (1986) reported CO₂ solubility as a function of temperature and salinity, and the solubility coefficients were 37.28–38.12 μM/L/torr at 26°C and 30–35 salinity (psu).²⁴ The hemolymph aco₂ of *P. fucata martensii* at 26.0°C (37.02 μM/L/torr) was similar to the coefficient reported by Cameron (1986). The mean value of hemolymph pK_{app} in this study was 5.995853, whereas the hemolymph pK_{app} values of other marine bivalves were 6.114 in *M. edulis* at 12°C,^{11,23} 6.07343 in *C. gigas* at 23°C,¹⁶ 6.2609 in *M. coruscus* at 24°C,¹⁵ 6.0641 in *M. nobilis* at 24°C,¹⁴ and 5.9987 in *P. margaritifera* at 26°C.¹³ The pK_{app} value is equal to the pH value at which it is most effective as a

buffer.²⁴ The most effective buffer pH in *P. fucata martensii* hemolymph seemed to be similar to the value in *P. margaritifera*, *C. gigas*, and *M. nobilis*.

In *in vitro* experiments, the pH decreased significantly with an increase in Pco₂ (Table 2). The hemolymph Pco₂ estimated using the correction equation was lower than that from the mean value of pK_{app}, nevertheless there was no statistical difference in [HCO₃⁻] (Table 3). αco₂ and pK_{app} vary with ionic strength and temperature,¹⁸ and the estimation of Pco₂ may be affected by temperature and salinity. Therefore, it is necessary to examine these parameters at various temperatures and salinities in order to increase the accuracy of the calculation of Pco₂ and to formulate the correction equation.

Using the mean values of hemolymph αco₂ and pK_{app} in this study, Pco₂ and [HCO₃⁻] of the hemolymph in adductor muscle of *P. fucata martensii* were calculated. The mean values of hemolymph Pco₂ and [HCO₃⁻] in this species were 1.69 torr and 2.14 mM/L, respectively. In other marine bivalves, the mean values of hemolymph Pco₂ and [HCO₃⁻] were 0.9 torr and 1.8 mM/L in *M. edulis* at 12°C,¹¹ 1.15 torr and 1.62 mM/L in *M. galloprovincialis* at 18°C,¹² 2.18 torr and 1.78 mM/L in *C. gigas* at 23°C,¹⁶ 1.55 torr and 1.44 mM/L in *M. nobilis* at 23°C,¹⁴ and 1.50 torr and 1.98 mM/L in *P. margaritifera* at 26°C.¹³ The hemolymph Pco₂ and [HCO₃⁻] of *P. fucata martensii* were approximately the same as those of *C. gigas* and *P. margaritifera*.

The non-bicarbonate buffer value (β_{NB}) is provided by the buffer capacity of the non-bicarbonate buffer system by mainly protein and protein residues,²⁵ and used to quantify the amount of buffering of the solution component. In *P. fucata martensii*, the hemolymph β_{NB} and T_{pro} were 0.967 slykes and 1.6 g/L, respectively. The hemolymph β_{NB} and T_{pro} of *M. edulis* were 0.4–0.622 slykes and 0.79 g/L. The hemolymph β_{NB} and T_{pro} in *P. fucata martensii* were higher than that in *M. edulis*. The high β_{NB} in *P. fucata martensii* was probably as a result of the high concentration of the total protein in the hemolymph. Although there are few reports on the relationship between non-bicarbonate buffer value and total protein concentration in the marine bivalves, β_{NB}

was 0.65 slykes in *M. galloprovincialis*,¹²⁾ 0.44 slykes in *M. coruscus*,¹⁵⁾ and 0.732 slykes in *C. gigas*.¹⁶⁾ The hemolymph β_{NB} of *P. fucata martensii* was higher than that of other marine bivalves. Therefore, *P. fucata martensii* should have a relatively high buffering capacity of the non-bicarbonate buffer system of the hemolymph in comparison with *M. edulis*, *M. galloprovincialis*, *M. coruscus*, and *C. gigas*. The changes in hemolymph pH in *P. fucata martensii* could be relatively smaller than that of Mytilidae and Ostreidae bivalves, even if environmental CO₂ concentrations increase.

References

- 1) Hayami I: Pteriidae. In: Okutani T (ed) Marine Mollusks in Japan. The second edition, Tokai University Press, Tokyo, 1179-1181 (2017)
- 2) Yamamoto K, Adachi S, Koube H: Direct method of measuring the filtration volume in the pearl oyster *Pinctada fucata martensii*. *J Nat Fish Univ*, **44**, 189-194 (1996)
- 3) Yamamoto K, Handa T, Nakamura M, Kitukawa K, Kita Y, Takimoto S, Nishikawa S: Effects of ozone-produced oxidants on respiration of the pearl oyster, *Pinctada fucata martensii*. *Aquaculture Sci*, **47**, 241-248 (1999)
- 4) Yamamoto K, Adachi S, Koube H: Effects of hypoxia on respiration in the pearl oyster *Pinctada fucata martensii*. *Aquaculture Sci*, **47**, 539-544 (1999)
- 5) Yamamoto K: Effects of water temperature on respiration in the pearl oyster *Pinctada fucata martensii*. *Aquaculture Sci*, **48**, 47-52 (2000)
- 6) Yamamoto K, Handa T, Matsubara T: Change on ventilation volume in the pearl oyster *Pinctada fucata martensii* associated with feeding. *Aquaculture Sci*, **58**, 447-451 (2010)
- 7) Yamamoto K, Handa T, Kondo M: Structure of the digestive diverticula of the pearl oyster *Pinctada fucata martensii*. *J Nat Fish Univ*, **52**, 31-43 (2004)
- 8) Yamamoto K, Handa T, Kondo M: Ctenidium structure of the pearl oyster *Pinctada fucata martensii* (Pteriidae: Pseudolamellibranchia: Bivalvia). *J Nat Fish Univ*, **57**, 81-110 (2008)
- 9) Handa T, Yamamoto K: The blood acid-base balance in the pearl oyster *Pinctada fucata martensii* after the surgery. *J Nat Fish Univ*, **60**, 57-61 (2011)
- 10) Handa T, Yamamoto K: The acid-base balance of the hemolymph in the pearl oyster *Pinctada fucata martensii* under normoxic conditions. *Aquaculture Sci*, **60**, 113-117 (2012)
- 11) Booth CE, McDonald DG, Walsh PJ: Acid-base balance in the sea mussel, *Mytilus edulis*. I. Effects of hypoxia and air-exposure on hemolymph acid-base status. *Mar Bio Lett*, **5**, 347-358 (1984)
- 12) Michelidis B, Ozounis C, Palaras A, Portner HO: Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Mar Ecol Prog Ser*, **293**, 109-118 (2005)
- 13) Handa T, Yamamoto K: Estimation of CO₂ partial pressure and bicarbonate concentration in the hemolymph of the black-lip pearl oyster *Pinctada margaritifera*. *J Nat Fish Univ*, **63**, 181-188 (2015)
- 14) Handa T and Yamamoto K: Estimation of CO₂ partial pressure and bicarbonate concentration in the hemolymph of the noble scallop *Mimachlamys nobilis*. *J Nat Fish Univ*, **64**, 188-194 (2016)
- 15) Handa T, Araki A, Yamamoto K: Acid-base balance of the hemolymph in hard-shelled mussel *Mytilus coruscus* in normoxic conditions. *J Nat Fish Univ*, **65**, 39-46 (2017)
- 16) Handa T, Araki A, Kawana K, Yamamoto K: Acid-base balance of hemolymph in Pacific oyster *Crassostrea gigas* in Normoxic conditions. *J Nat Fish Univ*, **66**, 103-110 (2018)
- 17) Handa T, Araki A, Yamamoto K: Oxygen and acid-base status of hemolymph in the densely lamellated oyster *Ostrea dendelamellosa* in Normoxic conditions. *J Nat Fish Univ*, **66**, 203-208 (2018)
- 18) Boutilier RG, Iwama GK, Heming TA, Randall DJ: The apparent pK of carbonic acid in rainbow trout blood plasma between 5 and 15°C. *Resp Physiol*, **61**, 237-254 (1985)
- 19) Yamamoto K, Adachi S, Tamura I, Aramizu T and

- Koube H: Effects of hypoxia and water temperature on ciliary movement of gills 5 bivalvia, *Mytilus edulis*, *Atrina pectinate*, *Pinctada fucata martensii*, *Chlamys nobilis* and *Crassostrea gigas*. *J Nat Fish Univ*, **44**, 137-142 (1996)
- 20) Yamamoto K, Handa T: Effect of hypoxia on oxygen uptake in the Pacific oyster *Crassostrea gigas*. *Aquaculture Sci*, **59**, 199-202 (2011)
- 21) Davenport HW: Fundamental equation. *In*: The ABC of acid–base chemistry 6th edition. University of Chicago Press, Chicago, 39-41 (1974)
- 22) Thomas RC: Intracellular pH. *In*: Hainsworth R (ed) Acid–base balance. Manchester University Press, United Kingdom, 50-74 (1986)
- 23) Walsh JP, McDonald DG, Booth CE: Acid–base balance in the sea mussel, *Mytilus edulis*. II. Effects of hypoxia and air-exposure on intracellular acid–base status. *Mar Biol Lett*, **5**, 359-369 (1984)
- 24) Cameron JN: The solubility of carbon dioxide as a function of temperature and salinity (Appendix table): *In*: Cameron JN (ed) Principles of physiological measurement. Academic Press, United Kingdom, 258-259 (1986)
- 25) Heisler N: Acid-base regulation: *In*: Evans DH (ed) The physiology of fishes. CRC Press, Florida, 343-378 (1993)

