

Estimation of α_{CO_2} and pKapp of hemolymph acid–base balance in *Mytilus coruscus* between 16°C and 28°C

Takeshi Handa[†] and Akira Araki

Abstract : The influence of temperature on the hemolymph CO_2 solubility coefficient (α_{CO_2}) and the apparent dissociation constant of carbonic acid (pKapp) in *Mytilus coruscus* was investigated. *Mytilus coruscus* hemolymph was equilibrated with standard CO_2 gas mixtures to obtain expressions for α_{CO_2} and pKapp as a function of temperature. The relationship between α_{CO_2} and temperature (T) is expressed as follows: $\alpha_{\text{CO}_2} = 138.247 - 11.253 \cdot T + 0.554 \cdot T^2 - 0.0140 \cdot T^3 + 0.000138 \cdot T^4$. The following relationship between pKapp and temperature was found: $\text{pKapp} = 6.6407 - 0.01589 \cdot T$. The parameter units are °C for T and $\mu\text{M/L/torr}$ for α_{CO_2} . These equations enable estimation of hemolymph α_{CO_2} and pKapp at arbitrary temperatures and simple calculation of Pco_2 and $[\text{HCO}_3^-]$.

Key words : *Mytilus coruscus*, hemolymph acid–base balance, CO_2 solubility (α_{CO_2}), apparent dissociation constant (pKapp), temperature effect, normoxia

Introduction

The hard-shelled mussel *Mytilus coruscus* is a Mytilidae bivalve classified under the order Mytiloida within the subclass Pteriomorphia¹⁾. In Japan, *M. coruscus* inhabits the rocky bottom of intertidal zones from Hokkaido to Kyushu¹⁾. Collected in the littoral zone, it is considered a premium seafood and is referred to as “Sendai-gai” in Miyagi prefecture, or “Seto-gai” in Yamaguchi prefecture.

Mytilus coruscus was previously studied in terms of larvae morphology²⁾, polymorphic microsatellite loci³⁾, microsatellite markers⁴⁾, influence of natural biofilm on the settlement mechanism⁵⁾, immune activities of hemocytes⁶⁾, hybrid molecular identification⁷⁾, light-responsive genes⁸⁾, and marine environment⁹⁾. In the context of respiratory physiology, the response to air exposure on hemolymph acid–base balance was studied, indicating a slow, partial compensation of metabolic acidosis¹⁰⁾. However, for *M. coruscus* hemolymph, there are few reports evaluating CO_2 dynamics and acid–base balance with respect to the influence of temperature on the CO_2 solubility coefficient (α_{CO_2}) and apparent

dissociation constant of carbonic acid (pKapp).

Under normoxic and normocapnic conditions, in blue mussel *M. edulis*, the hemolymph CO_2 partial pressure (Pco_2) was 0.9 torr¹¹⁾, for Akoya pearl oyster *Pinctada fucata martensii* it was 1.7–2.3 torr^{12,13)}, and 1.55 torr for the noble scallop *Mimachlamys nobilis*¹⁴⁾. As bivalves generally have very low Pco_2 values, the same was expected for Pco_2 in *M. coruscus*; however, direct Pco_2 measurement is difficult. Estimation of Pco_2 via the Henderson–Hasselbalch equation is often used in studying acid–base balance owing to its ease and accuracy¹⁵⁾. In the equation, α_{CO_2} and pKapp values are required for the experimental animal. As temperature influences α_{CO_2} and pKapp¹⁵⁾, it is an important factor in determining *M. coruscus* hemolymph α_{CO_2} and pKapp. Therefore, the effect of temperature on *M. coruscus* hemolymph α_{CO_2} and pKapp was investigated. For *M. coruscus*, a simple relationship between temperature, hemolymph α_{CO_2} , and pKapp allows Pco_2 calculation with ease, contributing to understanding the relationships between acid–base balance, respiratory physiology, and aquaculture environments.

Materials and Methods

Experimental animals and conditions

These experiments used 85 hard-shelled mussel *M. corsucus* (mean wet weight: 180 g). They were collected from the coast of the Seto Inland Sea in the eastern area of Yamaguchi prefecture. After cleaning the shell valves, the mussels were reared in water with seasonal variations (16-28°C) and fed cultivated phytoplankton¹⁶⁻¹⁸⁾. Twenty-four hours before hemolymph collection, the mussels were transferred to particle-free (> 0.45 µm) seawater. All experiments were conducted in seawater with a salinity of 29 psu, O₂ saturation 96%, pH 8.0, and a total CO₂ concentration of 1.5 mM/L.

Hemolymph collection

Adductor muscle hemolymph was collected anaerobically by direct puncture with a gas-tight microsyringe (Model 1750LTN, Hamilton Co.), collecting approximately 0.3-0.4 mL. The hemolymph was used for analyses of αCO_2 and pKapp in *in vitro* experiments.

Experimental protocols

Analysis of αCO_2 used hemolymph adjusted to pH2.5 by lactic acid (Wako Pure Chemical Industries, Ltd.) addition. The acidified sample was transferred to a tonometer flask and equilibrated with humidified standard CO₂ gas (CO₂, 5.0% or 15%; O₂, 20.9%; N₂ Balance) using an equilibrator (DEQ-1, Cameron Instruments) at experimental temperature. After equilibration, the sample Tco₂ was measured (n=41). The Pco₂ of the equilibrated sample was calculated using the known standard gas CO₂ concentration, barometric pressure, and water vapor pressure. The αCO_2 was calculated using the equation:

$$\alpha\text{CO}_2 = \text{Tco}_2 \cdot \text{Pco}_2^{-1}$$

For pKapp determination, the hemolymph was used immediately after collection for tonometry analysis. Hemolymph was equilibrated with humidified standard CO₂ mixes (CO₂, 0.2-5.0%; O₂, 20.9%; N₂ Balance) with the equilibrator at 16 °C (n=25), 22 °C (n=30), or 28 °C (n=30).

After equilibration, the pH and Tco₂ of the sample were measured. Using the sample pH, Tco₂, and αCO_2 , calculated from the above equation, pKapp was determined by rearrangement of the Henderson-Hasselbalch equation^{15,19)} as follows:

$$\text{pKapp} = \text{pH} - \log [(\text{Tco}_2 - \alpha\text{CO}_2 \cdot \text{Pco}_2) \cdot (\alpha\text{CO}_2 \cdot \text{Pco}_2)^{-1}]$$

where Pco₂ was calculated from the known CO₂ concentration of standard gases.

Hemolymph analysis

Tco₂ was measured using a total CO₂ analyzer (Capnicon 5, Cameron Instruments). The pH was measured using a blood gas meter (BGM200, Cameron Instruments) with pH glass and reference electrodes (E301, E351, Cameron Instruments). The pH electrodes were installed in a water jacket and maintained at experiment temperatures. Hemolymph Pco₂ and [HCO₃⁻] were calculated by rearranging the Henderson-Hasselbalch equation^{15,19)}. The obtained αCO_2 and pKapp were then used to calculate hemolymph Pco₂ from pH and Tco₂:

$$\text{Pco}_2 = \text{Tco}_2 \cdot [\alpha\text{CO}_2 \cdot (1 + 10^{(\text{pH} - \text{pKapp})})]^{-1}$$

The bicarbonate ion ([HCO₃⁻]) concentration was calculated from Tco₂, αCO_2 and Pco₂, or from αCO_2 , Pco₂, pH, and pKapp using the equations:

$$\begin{aligned} [\text{HCO}_3^-] &= \text{Tco}_2 - \alpha\text{CO}_2 \cdot \text{Pco}_2 \\ [\text{HCO}_3^-] &= \alpha\text{CO}_2 \cdot \text{Pco}_2 \cdot 10^{(\text{pH} - \text{pKapp})} \end{aligned}$$

To assess the relationship between hemolymph pH and [HCO₃⁻] in the experimental animals, the non-bicarbonate buffer values (β_{NB}) were calculated from the slope of the relational expression between pH and [HCO₃⁻].

Statistical analysis

The Kruskal-Wallis test was performed for changes in hemolymph sample properties and the calculated pKapp. Multiple comparison for all pairs used the Steel-Dwass

test. Statistically significant differences were set at $P < 0.05$. All analyses were carried out using the statistical software Kyplot 6.0 (KyensLab Inc., Japan).

Results and Discussion

The influence of temperature on *M. coruscus* hemolymph αCO_2 and pKapp was investigated, and the relationship clarified. The hemolymph αCO_2 of *M. coruscus* are shown in Fig. 1. Analysis of samples over 30°C used hemolymph collected from animals reared at 28°C, and sample analysis at 15°C used hemolymph collected from animals reared at 16°C. The mean αCO_2 are 29.1–54.3 $\mu\text{M}/\text{L}/\text{torr}$ between 15°C and 34°C. Cameron (1986) reported CO_2 solubility as a function of temperature and salinity, with solubility coefficients of 31.58–53.45 $\text{mM}/\text{L}/\text{torr}$ at a salinity of 30 between 14°C and 34°C²⁰. The obtained hemolymph αCO_2 reflected that reported by Cameron (1986)²⁰. Although the information with respect to this point is limited, the hemolymph αCO_2 decreased with increasing temperature in *M. coruscus*, which matched responses of other animals; hemolymph of disk abalone *Haliotis (Nardotis) discus discus* under 20°C²¹ and plasma of rainbow trout *Salmo gairdneri* under 15°C¹⁵. In the hemolymph of *M. coruscus*, a polynomial equation is fitted to the αCO_2 data (Fig. 1), with this hemolymph αCO_2 equation estimating αCO_2 within the temperature range. The relationship between αCO_2 and temperature is expressed as follows:

$$\alpha\text{CO}_2 = 138.2475 - 11.2533 \cdot T + 0.553901 \cdot T^2 - 0.01399 \cdot T^3 + 0.000138 \cdot T^4 \quad (R^2 = 0.9384)$$

where T is the temperature, and units used are $\mu\text{M}/\text{L}/\text{torr}$ for αCO_2 and °C for T.

Hemolymph pKapp at each temperature is shown with corresponding Pco_2 , pH, and Tco_2 (Tables 1-3). The mean hemolymph pKapp were 6.38068 at 16°C, 6.30234 at 22°C, and 6.18990 at 28°C. These mean values enable calculation of hemolymph Pco_2 and $[\text{HCO}_3^-]$ at each temperature.

Hemolymph properties and pKapp at known Pco_2 are shown in Tables 1-3. Hemolymph pH and pKapp significantly change with increasing Pco_2 at each

temperature ($P < 0.05$, Kruskal–Wallis test). A polynomial equation was fitted to the mean values of pKapp and pH, yielding a relationship between pKapp and pH as follows:

$$16^\circ\text{C} \quad \text{pKapp} = -32.557 + 12.885 \cdot \text{pH} - 1.265 \cdot \text{pH}^2 - 0.0316 \cdot \text{pH}^3 \quad (R^2 = 0.9983)$$

$$22^\circ\text{C} \quad \text{pKapp} = 144.863 - 64.994 \cdot \text{pH} + 10.106 \cdot \text{pH}^2 - 0.521 \cdot \text{pH}^3 \quad (R^2 = 0.9787)$$

$$28^\circ\text{C} \quad \text{pKapp} = 537.058 - 241.340 \cdot \text{pH} + 36.558 \cdot \text{pH}^2 - 1.845 \cdot \text{pH}^3 \quad (R^2 = 0.8827)$$

These equations enable pKapp estimation for each temperature, with hemolymph Pco_2 and $[\text{HCO}_3^-]$ calculation by Henderson–Hasselbalch equation rearrangement.

The distribution of pKapp and corresponding pH is shown for each temperature in Fig. 2. The distribution of pKapp at 28°C was different to that at 16°C and 22°C ($P < 0.05$, Steel–Dwass test). There was no significant difference in pH distribution. The linear regression was fitted to the mean values of pKapp and temperature (Fig. 3), and the relationship between pKapp and temperature expressed as follows:

$$\text{pKapp} = 6.6407 - 0.01589 \cdot T \quad (R^2 = 0.9894)$$

where T is temperature in °C.

This equation estimates hemolymph pKapp at arbitrary temperatures (between 16–28°C). Thus, the hemolymph Pco_2 and $[\text{HCO}_3^-]$ may be calculated for a range of physiologically relevant temperatures.

Hemolymph pH and calculated $[\text{HCO}_3^-]$ with the Pco_2 of standard gases are listed in Table 4. The non-bicarbonate buffer values (β_{NB}), obtained as a regression coefficient relating pH and $[\text{HCO}_3^-]$, were 0.40 slykes at 16°C, 0.47 slykes at 22°C, and 0.29 slykes at 28°C. The non-bicarbonate buffer value was determined by the buffer capacity of the non-bicarbonate buffer system (for example, protein buffer system), and used to quantify buffering of the solution component^{22,23}. Therefore, the

hemolymph of *M. coruscus* has a greater buffer capacity at 16°C and 20°C than at 28°C. For other mussels, hemolymph β_{NB} in *M. edulis* was 0.4 slykes at 12°C¹¹⁾, and 0.65 slykes in *M. galloprovincialis* at 18°C²⁴⁾. The hemolymph β_{NB} of *M. coruscus* reflected other mussels, with the hemolymph buffer capacity of the non-bicarbonate buffer system expected to reflect the Mitilid species.

In *M. coruscus*, the effect of temperature on hemolymph α_{CO_2} and pKapp were investigated, values required to calculate P_{CO_2} and $[HCO_3^-]$. As temperature strongly affects hemolymph α_{CO_2} and pKapp in *M. coruscus*, the proposed equations enable estimation of hemolymph α_{CO_2}

and pKapp at arbitrary temperatures and calculation of both P_{CO_2} and $[HCO_3^-]$ with relative ease. Boutilier et al. (1985) described variations in plasma α_{CO_2} and pKapp for rainbow trout in response to temperature and ionic strength, with pKapp changes influenced by plasma pH¹⁵⁾. If it is possible to similarly discuss acid-base balance of the fish and the mussel, further exploration of the acid-base balance under resting conditions and relationships among temperature, pH and pKapp of *M. coruscus* is necessary.

Table 1. Mean values of measured pH, total CO₂ content (Tco₂) and calculated apparent dissociation constant of carbonic acid (pKapp) of *Mytilus coruscus* hemolymph with known Pco₂ standard gases at 16°C

Standard gas		Hemolymph			
CO ₂	Pco ₂	pH	Tco ₂	pKapp	n
(%)	(torr)		(mM/L)		
0.2	1.5	7.538	1.67	6.2197382	5
0.5	3.7	7.305	1.86	6.3690807	5
1.0	7.4	7.119	2.13	6.4608385	5
2.0	14.7	6.794	2.29	6.4940144	5
5.0	36.8	6.382	3.95	6.3597565	5

Water temperature, 16.0°C; Mean value of pKapp, 6.3806857.

Table 2. Mean values of measured pH, total CO₂ content (Tco₂) and calculated apparent dissociation constant of carbonic acid (pKapp) of *Mytilus corsucus* hemolymph with known Pco₂ standard gases at 22°C

Standard gas		Hemolymph			
CO ₂	Pco ₂	pH	Tco ₂	pKapp	n
(%)	(torr)		(mM/L)		
0.2	1.5	7.485	1.56	6.117095	6
0.5	3.7	7.297	1.66	6.331092	6
1.0	7.5	7.088	1.92	6.383487	6
2.0	14.9	6.777	2.10	6.412913	6
5.0	37.3	6.355	3.57	6.267141	6

Water temperature, 21.8°C; Mean value of pKapp, 6.3023457.

Table 3. Mean values of measured pH, total CO_2 content (Tco_2) and calculated apparent dissociation constant of carbonic acid (pKapp) of *Mytilus coruscus* hemolymph with known Pco_2 standard gases at 28°C


Standard gas		Hemolymph			
CO_2	Pco_2	pH	Tco_2	pKapp	n
(%)	(torr)		(mM/L)		
0.2	1.5	7.196	1.02	5.924225	6
0.5	3.7	7.091	1.14	6.197768	6
1.0	7.3	6.830	1.29	6.227287	6
2.0	14.6	6.589	1.50	6.307682	6
5.0	36.6	6.277	2.53	6.292557	6

Water temperature, 28.0°C ; Mean value of pKapp, 6.1899038.

Table 4. Mean values of hemolymph pH and bicarbonate concentration ($[\text{HCO}_3^-]$) of *Mytilus coruscus* with known Pco_2 standard gases at 16-28°C

Standard gas	16°C		22°C		28°C	
	pH	$[\text{HCO}_3^-]$	pH	$[\text{HCO}_3^-]$	pH	$[\text{HCO}_3^-]$
(%)		(mM/L)		(mM/L)		(mM/L)
0.2	7.538	1.59	7.485	1.49	7.196	0.97
0.5	7.305	1.66	7.297	1.50	7.091	1.01
1.0	7.119	1.75	7.088	1.60	6.830	1.03
5.0	6.382	2.04	6.355	1.99	6.277	1.24

The non-bicarbonate buffer value (β_{NB}), 0.40 slykes at 16°C ; 0.47 slykes at 22°C ; 0.29 slykes at 28°C.

**Fig. 1.** Influence of temperature on CO_2 solubility coefficient (α_{CO_2}) for *Mytilus coruscus* hemolymph. Data is the calculation value. The dotted line is fitted to the data and the equation: $\alpha_{\text{CO}_2} = 138.2475 - 11.2533 \cdot T + 0.553901 \cdot T^2 - 0.01399 \cdot T^3 + 0.000138 \cdot T^4$ ($R^2 = 0.9384$, $n=41$).




Fig. 2. The pKapp distribution and corresponding pH of the hemolymph in *Mytilus corsucus* at 16°C, 22°C and 28°C. Data is the calculation value. Open circle, 16°C (n=25); solid circle, 22°C (n=30); open square, 28°C (n=30). The pKapp distribution at 28°C was different to that at 16°C and 22°C ($P < 0.05$, Steel-Dwass multiple comparison test).




Fig. 3. Influence of temperature on the apparent carbonic acid dissociation constant (pKapp) of *Mytilus corsucus* hemolymph between 16°C and 28°C. Data are Mean \pm standard error. The dotted line is fitted to the data and the equation: $\text{pKapp} = 6.6407 - 0.01589 \cdot T$ ($R^2 = 0.9894$).

Acknowledgments

We would like to express our sincere gratitude to Dr. Ken-ichi Yamamoto, Professor Emeritus, for securing the experimental animals for this study.

References

- 1) Kurozumi T: Mytiloida. In: Okutani T (ed) Marine mollusks in Japan (2nd edition), Tokai University Press, Kanagawa, 1172-1179 (2017)
- 2) Semenikhina OY, Kolotukhina NK, Evseev GA: Morphology of larvae of the family Mytilidae (Bivalvia) from the north-western part of the Sea of Japan. *Journal of the Marine Biological Association of the United Kingdom*, **88**, 331-339 (2008)
- 3) Xu TJ, Sun YN, Yuan YT, Liao Z, Wang RX: Isolation and characterization of polymorphic microsatellite loci in the hard-shelled mussel, *Mytilus coruscus* (Mytilidae). *Genetics Molecular Research*, **9**, 1338-1391 (2010)
- 4) An HS, Lee JW: Development of microsatellite markers for the Korean mussel, *Mytilus coruscus* (Mytilidae) using next-generation sequencing. *International Journal of Molecular Science*, **13**, 10583-10593 (2012)
- 5) Yand JL, Li X, Liang X, Bao WY, Shen HD, Li JL: Effects of natural biofilms on settlement of plantigrades of the mussel *Mytilus coruscus*. *Aquaculture*, **425**, 228-233 (2014)
- 6) Yang HS, Hong HK, Donaghy L, Noh CH, Park HS, Kim DS, Choi KS: Morphology and immune-related activities of hemocytes of the mussel *Mytilus coruscus* (Gould, 1861) from East Sea of Korea. *Ocean Science Journal*, **50**, 77-85 (2015)
- 7) Zhang W, Li R, Chen X, Wang C, Gu Z, Mu C, Song W, Zhan P, Huang J: Molecular identification reveals hybrids of *Mytilus coruscus* x *Mytilus galloprovincialis* in mussel hatcheries of China. *Aquaculture International*, **28**, 85-93 (2020)
- 8) Xu M, Li J, Guo B, Qi P, Ye Y, Yan X: Identification and characterization of light-responsive genes in pre-settlement eyed veligers of *Mytilus coruscus*. *Aquaculture Reports*, **33**, 101768 (2023)
- 9) Qian J, Deng F, Shumway SE, Hu M, Wang Y: The thick-shell mussel *Mytilus coruscus*: Ecology, physiology, and aquaculture. *Aquaculture*, **580**, 740350 (2024)
- 10) Handa T, Araki A, Yamamoto K: Effect of air exposure on acid-base balance of hemolymph in hard-shelled mussel *Mytilus coruscus*. *Journal of National Fisheries University*, **68**, 65-70 (2020)
- 11) Booth CE, McDonald DG, Walsh PJ: Acid-base balance in the sea mussel, *Mytilus edulis*. I. Effects of hypoxia and air-exposure on hemolymph acid-base status. *Marine Biology Letters*, **5**, 347-358 (1984)
- 12) Handa T, Yamamoto K: The blood acid-base balance in the pearl oyster, *Pinctada fucata martensii*, after the surgery. *Journal of National Fisheries University*, **60**, 57-61 (2011)
- 13) Handa T, Yamamoto K: The acid-base balance of the hemolymph in the pearl oyster *Pinctada fucata martensii* under normoxic conditions. *Aquaculture Science*, **60**, 113-117 (2012)
- 14) Handa T, Yamamoto K: Estimation of CO₂ partial pressure and bicarbonate concentration in the hemolymph of the noble scallop *Mimachlamys nobilis*. *Journal of National Fisheries University*, **64**, 188-194 (2016)
- 15) Boutilier RG, Iwama GK, Heming TA, Randall DJ: The apparent pK of carbonic acid in rainbow trout blood plasma between 5 and 15°C. *Respiratory Physiology*, **61**, 237-254 (1985)
- 16) Yamamoto K, Adachi S, Tamura I, Aramizu T, Koube H: Effects of hypoxia and water temperature on ciliary movement of gills 5 bivalvia, *Mytilus edulis*, *Atrina pectinate*, *Pinctada fucata martensii*, *Chlamys nobilis* and *Crassostrea gigas*. *Journal of National Fisheries University*, **44**, 137-142 (1996)
- 17) Yamamoto K, Handa T, Nakamura M, Kitukawa K, Kita Y, Takimoto S, Nishikawa S: Effects of ozone-produced oxidants on respiration of the pearl oyster, *Pinctada fucata martensii*. *Aquaculture Science*, **47**, 241-248 (1999)
- 18) Yamamoto K, Handa T: Effect of hypoxia on oxygen uptake in the Pacific oyster *Crassostrea gigas*. *Aquaculture Science*, **59**, 199-202, (2011)
- 19) Davenport HW: Fundamental equation. In: Davenport

- HW (ed) The ABC of acid-base chemistry (6th edition), University of Chicago Press, Chicago, 39-41 (1974)
- 20) Cameron JN: The Solubility of carbon dioxide as a function of temperature and salinity (appendix table). *In*: Cameron JN (ed) Principles of physiological measurement. Academic Press, London, 258-259 (1986)
- 21) Handa T, Araki A: Estimation of hemolymph αCO_2 and pKapp in disk abalone *Haliotis (Nordotis) discus discus* between 10°C and 20°C. *Journal of National Fisheries University*, **72**, 103-111 (2024)
- 22) Heisler N: Acid-base regulation, interrelationships between gaseous and ionic exchange. *In*: Boutilier RG (ed) Vertebrate gas exchange, comparative & environmental physiology 6. Springer-Verlag, Berlin & Heidelberg, 211-251 (1990)
- 23) Claiborne JB: Acid-base regulation. *In*: Evans DH (ed) The physiology of fishes (2nd edition). CRC Press LLC, Florida, 177-198 (1998)
- 24) Michaelidis B, Ouzounis C, Paleras A, Pörtner HO: Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Marine Ecology Progress Series*, **293**, 109-118 (2005)

水温16℃から28℃のイガイにおける ヘモリンパ液の二酸化炭素溶解度と炭酸解離恒数の推定

半田岳志, 荒木 晶

和文要旨: イガイの呼吸機能, 特に酸塩基平衡を評価するため, イガイ閉殻筋から採取したヘモリンパ液を用いて, 二酸化炭素溶解度 (α_{CO_2}) と炭酸解離恒数 (pKapp) に及ぼす温度の影響について調査した。各実験温度において, イガイのヘモリンパ液を二酸化炭素標準ガスと平衡させ, pHと全炭酸含量を測定し, 温度 (T) と α_{CO_2} あるいはpKappとの関係を分析したところ, 以下の関係式を得た。 $\alpha_{\text{CO}_2} = 138.247 - 11.253 \cdot T + 0.554 \cdot T^2 - 0.0140 \cdot T^3 + 0.000138 \cdot T^4$, $\text{pKapp} = 6.6407 - 0.01589 \cdot T$ (α_{CO_2} : $\mu\text{M/L/torr}$; T: $^{\circ}\text{C}$)。これらの式により, 任意の温度でイガイのヘモリンパ液における α_{CO_2} とpKappの推定が可能となった。これら推定値を用いれば, 微量なヘモリンパ液の二酸化炭素分圧や炭酸水素イオン濃度を任意の温度で把握することができるだろう。